



Detroit R&D, Inc
2727 Second Ave. Suite 4113
Detroit, MI 48201
Phone: (313) 961-1606
Email: info@DetroitRandD.com
Web: www.DetroitRandD.com

Oxidative Stress ELISA (8-isoprostane) Kit

(iPF2 α -III, 15-isoprostane F2t, 8-epi-prostaglandin F2 α , 8-iso-prostaglandin F2 α)

Cat # 8iso1: ELISA kit for measuring 8-isoprostane in biological / wastewater samples

Cat # 8iso1R: 8-isoprostane ELISA Kit with removable strips

This competitive ELISA kit is for determination of 8-isoprostane levels in biological samples. A typical standard curve of the ELISA is shown in the bottom left corner.


The isoprostanes are a family of eicosanoids of non-enzymatic origin produced by the random oxidation of tissue phospholipids by oxygen radicals⁴. 8-epi-prostaglandin F2 α is an *in vivo* index of oxidative injury and associated with increased coronary artery calcification (CAC). A recent study using the 8-isoprostane ELISA revealed that rat hepatic 8-isoprostane levels increased in high fat (HF) diet-induced non-alcoholic fatty liver disease (NAFLD)¹⁸. Interestingly, telmisartan, an angiotensin II receptor antagonist (hypertension drug), significantly decreased the liver 8-isoprostane levels. Urinary excretion of 8-isoprostane increased in mice by aldosterone infusion-induced renal fibrosis. The level of 8-isoprostane excretion markedly decreased in osteopontin (involved in tumor promotion)-knock-out mice, suggesting that osteopontin is a therapeutic target for oxidative stress-induced kidney injury¹⁷. Levels of both renal NADPH oxidase subunit mRNA expression and urinary 8-isoprostane were increased in streptozotocin-induced diabetes mellitus (DM)¹⁶. 8-isoprostane has been examined as an exercise-induced muscle damage biomarker¹⁵ and as a sewage biomarker for assessing community health¹⁵. Yamamoto et al. showed that the use of AST-120 was related to improvement of oxidative stress (decrease in 8-isoprostane levels) in patients undergoing maintenance haemodialysis treatment¹³. *In utero* diesel exhaust exposure of hyperlipidemic mice did not influence later life redox homeostasis¹¹. A strong association between tobacco smoking and urinary 8-isoprostane levels was found in a large German study of older adult smokers^{4,6,7,8,12}. Soleus muscle from T2D-induced rats fed a high fat diet exhibited higher levels of 8-isoprostane in conjunction with increased mt DNA copy number and 8-OHdG⁹. Hyperoxia increased 8-isoprostane in lung of wild type mice but not in soluble epoxide hydrolase knockout mice¹⁰. 8-Isoprostane levels were related with cardiovascular disease risk and environmental estrogen exposure^{1,3}. Diluted urine or cell culture media without extraction are suitable for the ELISA.

Each kit for triplicate analyses of up to 24 samples contains one 96 well plate, one tube of 8-isoprostane standard, one tube of 8-isoprostane-conjugated horseradish peroxidase (HRP), and buffers for sample and HRP dilutions, and plate washing.

Related ELISA kits:

Environmental Estrogen BPA and BPS
ELISA kits

Air Pollution 2-Naphthol (2-NAP) ELISA
kit



1. Santos JM, Zhu J, McPike A, Dechant J, Demirci J. Biomonitoring Endocrine Disruptor Chemicals in Human Milk: A New Approach for Understanding Cardiovascular Disease Risk in Women Living in Environmental Justice Areas. The SOT abstract, #3058. 2025.
2. Pratt K, Hilty A, Jacob P, Schick SF. Respiratory Exposure to Thirdhand Cigarette Smoke Increases Concentrations of Urinary Metabolites of Nicotine. *Nicotine and Tobacco Research* 25.8. 2023.
3. Riiffee JNJ, Wade M, Sine T, Griffith C, Benite-Ribeiro S, Haney A, Santos JM. The link of environmental estrogens exposure to oxidative stress and their association with insulin- and exercise-induced glucose uptake. *Obesity Medicine* 42. 2023.
4. Gào X, Xuan Y, Benner A, Anusruti A, Brenner H, Schöttker, B. Nitric Oxide Metabolites and Lung Cancer Incidence: A Matched Case-Control Study Nested in the ESTHER Cohort. *Oxidative Medicine and Cellular Longevity* Volume 2019, Article ID 6470950.
5. Kawasaki K, Kondoh E, Chigusa Y, Kawamura Y, Mogami H, Takeda S, Horie A, Baba T, Matsumura N, Mandai M, Konishi I. Metabolomic profiles of placenta in preeclampsia: Antioxidant effect of magnesium sulfate on trophoblasts in early-onset preeclampsia. *Hypertension* 73: 671-679. 2019.
6. Gào X, Zhang Y, Burwinkel B, Xuan Y, Holleccek B, Brenner H, Schöttker B. The associations of DNA methylation alterations in oxidative stress-related genes with cancer incidence and mortality outcomes: a population-based cohort study. *Clinical Epigenetics* 11: 14-22. 2019.

7. Gao, X, Gào X, Zhang Y, Holleczeck B, Schöttker B Brenner H. Oxidative stress and epigenetic mortality risk score: associations with all-cause mortality among elderly people. *Eur. J. Epidemiology* 34: 451-462. 2019.
8. Gào X, Brenner H, Holleczeck B, Cuk K, Zhang Y, Anusruti A, Xuan Y, Xu Y, Schöttker B. Urinary isoprostane levels and occurrence of lung, colorectal, prostate, breast and overall cancer: Results from a large, population-based cohort study with 14 years of follow-up. *Free Radical Biology and Medicine* 123: 20-26. 2018.
9. Santos, JM et al. The role of mitochondrial DNA damage at skeletal muscle oxidative stress on the development of type 2 diabetes. *Molecular and Cellular Biochemistry* doi.org/10.1007/s11010-018-3361-5. 2018.
10. Liu, LP et al. Deletion of soluble epoxide hydrolase attenuates mice hyperoxic acute lung injury. *BMC Anesthesiology* doi.org/10.1186/s12871-018-0490-z. 2018.
11. Harrigan et al. In utero exposure of hyperlipidemic mice to diesel exhaust: Lack of effects on atherosclerosis in adult offspring fed a regular chow diet. *Cardiovasc Toxicol.* DOI:10.1007/s12012-017-9399, 2017.
12. Gao et. al. Associations of self-reported smoking, cotinine levels and epigenetic smoking indicators with oxidative stress among older adults: a population-based study. *Genetic Epidemiology* 32: 443-456, 2017.
13. Yamamoto et al. Continuous reduction of protein-bound uraemic toxins with improved oxidative stress by using the oral charcoal adsorbent AST-120 in haemodialysis patients. *Sci. Rep.* 5, 14381; doi:1 0.1038/srep14381 2015.
14. Santos JM, Jurban M, Kim H. Could sewage epidemiology be a strategy to assess lifestyle and wellness of a large scale population? *Medical Hypothesis* 85: 408-411, 2015.
15. Kanda et al. Evaluation of serum leaking enzymes and investigation into new biomarkers for exercise-induced muscle damage. *EIR* 20: 39-54, 2014.
16. Nagao T, Okura T et al. Osteopontin plays a critical role in interstitial fibrosis but not glomerular sclerosis in diabetic nephropathy. *Nephron Extra* 2, 87-103, 2012.
17. Irita et al. Osteopontin deficiency protects against aldosterone-induced inflammation, oxidative stress, and interstitial fibrosis in the kidney. *Am. J. Physiol. Renal Physiol.* 301, F833-844, 2011.
18. Sookoian et al. High fat diet-induced liver steatosis promotes an increase in liver mitochondrial biogenesis in response to hypoxia. *J. Cell. Mol. Med.* 15, 1329-1338, 2011.